**REMARKS** 

**Specification Objections:** 

The specification was objected to because a less than complete description of the

plant was presented. Accordingly, the specification has been amended along the lines of

the examiner's suggestions to more clearly and completely describe the plant. It is

respectfully requested that the specification objection be withdrawn.

Rejection Under 35 U.S.C. 112:

The claim was rejected under 35 USC 112 as not being supported by a clear

botanical description. Accordingly, the botanical description has been amended to more

clearly describe the plant. It is respectfully requested that the 112 rejection be

withdrawn.

**Conclusion:** 

The Examiner's attention to each of the parts of the patent application is greatly

appreciated.

The claimed plant is now believed to be in condition for allowance.

Respectfully submitted,

PLANTHAVEN, INC.

Geoffrey Needham, President

Tel: (805) 569 - 6498



### EUPHORBIA PLANT NAMED 'FIRST BLUSH'

-GENUS: EUPHORBIA

5

-SPECIES: epithymoides

DENOMINATION: FIRST BLUSH

10

15

20

25

#### **BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of variegated cushion spurge, a hardy perennial that is grown for use as an ornamental shrub for the landscape. The new invention is known botanically as *Euphorbia* and will be referred to hereinafter by the cultivar name 'FIRST BLUSHFirst Blush'.

'FIRST BLUSHirst Blush' is a sport that was discovered by the inventor in 1996 in a cultivated area of Germantown, Wisconsin. 'FIRST BLUSHirst Blush' was discovered as a variegated branch on an unnamed individual plant of Euphorbia epithymoides among a production block grown from seed. 'FIRST BLUSHirst Blush' was selected by the inventor based on its variegated foliage and variegated bracts. The foliage is pale green with cream-white margins. In spring the upper foliage nearest the sun is tinted rose-pink and then turns a more pronounced rose-red color in fall. Whorls of bracts are pale green with cream-white margins on stems that are often tinted red. 'FIRST BLUSHirst Blush' exhibits low vigor resulting in a less invasive plant than the species. The variegation makes 'FIRST BLUSHirst Blush' a showier variety than other Euphorbia known to the inventor.

The first asexual reproduction of 'FIRST BLUSHirst Blush' was carried out in a cultivated area of Germantown, Wisconsin by the inventor. The method used was stem

cuttings. Since that time subsequent generations have been determined stable and true to type.

#### SUMMARY OF THE INVENTION

5

10

15

20

The following traits have been repeatedly observed and represent the distinguishing characteristics of the new *Euphorbia* cultivar 'FIRST BLUSHirst Blush'. These traits in combination distinguish 'FIRST BLUSHirst Blush' from all other existing varieties of *Euphorbia* known to the inventor. 'FIRST BLUSHirst Blush' has not been tested under all possible conditions and phenotypic differences may be observed with variations in environmental, climatic, and cultural conditions, however, without any variance in genotype.

- 1. Euphorbia 'FIRST BLUSHirst Blush' exhibits a compact, clumping habit.
- 2. *Euphorbia* 'FIRST BLUSHFirst Blush' exhibits variegated foliage with pale green centers and cream-white margins with slightly crinkled edges.
- 3. *Euphorbia* 'FIRST BLUSHirst Blush' exhibits whorls of variegated pale green and cream-white bracts.
- 4. The edges of the upper foliage on *Euphorbia* 'F<u>IRST BLUSH</u>irst Blush' turn rose-pink in spring and deepen in color to rose-red in fall.
- 5. *Euphorbia* 'FIRST BLUSHirst Blush' exhibits low vigor and is less invasive than the species.
- 6. Euphorbia 'FIRST BLUSHirst Blush' is hardy to USDA Zone 4b.
- 7. Euphorbia 'FIRST BLUSHirst Blush' is winter dormant.
- 8. Euphorbia 'FIRST BLUSHirst Blush' exhibits yellow flowers in spring.

25

30

#### BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying drawings illustrate the distinguishing traits of the new *Euphorbia* cultivar '<u>FIRST BLUSHFirst Blush</u>'. The drawings were made in spring and summer of plants grown out-of-doors in a cultivated area of Germantown, Wisconsin.

The plants were in one-litre containers at the time. The drawing on sheet 1 illustrates the variegated foliage with rose-pink edges. The drawing on sheet 2 illustrates the plant in bloom. The drawings were made using conventional techniques and although flower and foliage colors may appear different from actual colors due to light reflectance, they are as accurate as possible by conventional photography.

#### BOTANICAL DESCRIPTION OF THE PLANT

The following is a detailed botanical description of the new *Euphorbia* cultivar

'FIRST BLUSHirst Blush'. Observations, measurements, values and comparisons were collected in Arroyo Grande, California from plants grown out-of-doors in one-litre containers. The observed plants were approximately nine to ten months old. The botanical data was collected as the plants grew back and matured after winter die-down, the plants having been started as cuttings in the late summer of the previous year. Color determinations are made in accordance with the 2001 Royal Horticultural Society Colour Chart from London England, except where general color terms of ordinary dictionary significance are used.

Botanical classification: Euphorbia 'FIRST BLUSHirst Blush'.

Species: *eEpithymoides*.

5

20 Common name: Variegated cushion spurge.

Parentage: Euphorbia 'FIRST BLUSHirst Blush' is a sport of the following parent plant:

Parent plantarent: An individual unnamed plant of Euphorbia epithymoides.

Propagation method: Division and cuttings.

Rooting habit: Fibrous.

25 Time to develop roots: 2-3 weeks are needed for initial cuttings to develop roots.

Crop time: 8 - 12 months are needed to produce a one-litre container from a rooted cutting.

Growth habit: Compact, clumping habit.

Use: Ornamental shrub for rock gardens and perennial borders.

30 Type: Perennial.

Height of plant: 18 cm. in height in a one-litre container.

Width of plant: 27 cm. in width in a one-litre container.

Cultural requirements: Plant in well-draining soil and full sunlight with moderate water.

Needs filtered shade in hot climates.

5 Hardiness: Hardy to USDA Zone 4b.

Seasonal traits: Yellow flowering in spring and <u>rose</u> tinted foliage <u>from in spring to</u> falland <u>summer</u>.

<u>ToxicityOther</u>: When cut or bruised plant exudes a-milky white latex that can be toxic and may elicit

10 dermal irritation.

Disease and pest resistance and susceptibility: Not resistant nor particularly susceptible to any diseases or pests known to the inventor.

Stem:

Branching habit: Basal branching.

15 Internode dimensions: 5 mm. between nodes.

Stem color: A combination of 138B and 61A.

Stem shape: Cylindrical.

Stem diameter: 3 mm. in diameter.

Stem length: 12 cm. in length.

Stem surface: Heavily pubescent.

Color of pubescence: 156D.

Foliage:

Type: Deciduous.

Leaf arrangement: Spiral.

Internode dimensions: 5 mm. between nodes.

Leaf division: Simple.

Leaf shape: Oblong elliptic.

Leaf length: 2 cm. in length.

Leaf width: 1 cm. in width.

Leaf apex: Acute.

Leaf base: Rounded.

Leaf venation pattern: Visible mid-vein.

5 Vein color: 138D.

Margin: Entire and revolute.

Leaf surface (abaxial surface): Pubescent.

Leaf surface (adaxial surface): Pubescent.

Leaf texture: Slightly crisped.

10 Leaf attachment: Petiole.

Petiole dimensions: 2.50 cm. in length and 1.25 mm. in width.

Petiole color: 138B.

Petiole surface: Glabrous.

Leaf color in spring (adaxial and abaxial surface): Centers are N138D and margins

are 158B with edges tinted 63C.

Leaf color in fall (adaxial and abaxial surface): Centers are N138D and margins are 158B with edges tinted 58A.

Bracts:

20

Bract color (adaxial surface): Center is N138D, margin is 158C and edge is 67A.

Bract color (abaxial surface): Center is 138C, margin is 158D and edge is 67D.

Number of bracts: Approximately 17 in number.

Bract arrangement: Whorled.

Fused or unfused: Unfused.

Bract dimensions: Range in size from 1.50 mm. in length to 6 mm. in length and

25 1.25 mm. in width to 5 mm. in width.

Bract surfaces (adaxial and abaxial surfaces): Pubescent.

Bract shape: Oblong.

Bract apex: Acute.

Bract base: Rounded.

30 Bract margins: Entire.

| <u>C</u>  | <u>Cyathia</u> Flower:  |
|-----------|---|
|           | Arrangement: Terminal cyathium.   |
|           | Number of cyathia: A range of 25-30 cyathium per 1-litre container.         |
|           | Dimensions of cyathia: 4 mm. in height and 6 mm. in width.                  |
|           | Cyathia color (adaxial and adaxial surfaces): Colors 11C, 138A, and 63C are |
| <u>i1</u> | ndividually present.  |
|           | Petals: Apetalous.  |
|           | Sepals: Asepalous.  |
|           | Inflorescence: Cyathuium.   |
|           | Flowering season: Mid sSpring to summer.                                    |
| _         | Lastingness of flower on plant: 5-10 days.                                  |
|           | Flower aspect: Facing upward.   |
|           | Fragrance: Absent.  |
|           | Peduncle color: 138B.   |
|           | Peduncle length: 1 cm. in length.   |
|           | Peduncle width: 1 mm. in width.   |
|           | Peduncle shape: Cylindrical.  |
|           | Peduncle surface: Glabrous.   |
|           | Shape of petaloid appendages floral leaves: Oblong elliptic.                |
|           | Color of petaloid appendages floral leaves: 11A.                            |
|           | Dimensions of petaloid appendages: 7 mm. in width and 14 mm. in length.     |
|           | Petaloid appendage surface (adaxial and abaxial surfaces): Glabrous.        |
| _         | Number of petaloid appendages: Six in number.                               |
|           | Petaloid appendage margins: Entire and sinuate.                             |
|           | Petaloid appendage apex: Rounded.   |
| ••••      | Petaloid appendage base: Attenuate.   |
|           | Reproductive organs:  |
|           | Stamen: <u>Six</u> One in number.   |
|           | Stamen color: 161A.   |
| _         | Stamen length: 1 mm. in length.   |

|    | Stamen form: Jointed filament.                               |
|----|--|
|    | Anther color: 161A.  |
|    | Anther dimensions: 0.25 mm. in width and 0.25 mm. in length. |
|    | Pollen amount: Low amount.                                   |
| 5  | Pollen color: 161A.  |
|    | Pistil: One in number.                                       |
|    | Pistil length: 1.50 mm. in length.                           |
|    | Pistil color: 11A.   |
|    | Styles: Three in number joined in a column.                  |
| 10 | Length of styles: 0.50 mm. in length.                        |
|    | Style color: 11A.  |
|    | Stigmas: Bifid.  |
|    | Stigma dimensions: 1 mm. in length and 1 mm. in width.       |
|    | Stigma color: 11A.   |
| 15 | Ovary: No ovary observed to date.                            |
|    | Seed:  |
|    | No sSeed production has not been observed to date.           |

# CLAIM

| A new and distinct cultivar          | of Euphorbia plant named | 'F <u>IRST</u> | <u>BLUSH</u> irst |
|--------------------------------------|--------------------------|----------------|-------------------|
| Blush' as described and illustrated. |                          |                |                   |

## **ABSTRACT**

A new cultivar of *Euphorbia* named 'FIRST BLUSHirst Blush' that is distinguished by a compact clumping habit, winter dormancy, showy variegated pale green foliage with rose colored tints, and yellow flowers in spring. In combination these traits set 'FIRST BLUSHirst Blush' apart from all other existing varieties of *Euphorbia* known to the inventor.